

DRAFT PRESCRIBING GUIDANCE PRESCRIBING CANNABIS MEDICINES FOR ANOREXIA/CACHEXIA IN ADVANCED CANCER

Introduction

The Australian Government's [Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia](#)¹ has identified little evidence for symptom benefit from cannabinoid medicines in palliative care patients with anorexia-cachexia syndrome.

Evidence-based care using proven interventions should be offered prior to cannabis medicines.

Current clinical trials

- **Medicinal Cannabis for Anorexia in Advanced Cancer**

A NSW Government funded [trial](#) to assess whether medicinal cannabis products can enhance the quality of life for adults with advanced cancer, particularly by improving appetite and appetite-related symptoms. See the [Australian New Zealand Clinical Trials Registry](#) for further information on the study.

Up to date information can be obtained from the [NSW Cannabis Medicines Advisory Service](#) by [email](#) or by telephone (02) 4923 6200.

Use of evidence-based therapies for cachexia-anorexia syndrome

Evidence-based management options in accordance with clinical practice guidelines, where available, must be offered to patients experiencing cachexia-anorexia syndrome in advanced cancer prior to considering the therapeutic use of cannabis medicines.

[Caresearch](#)² – the Palliative Care Knowledge Network – has resources available to assist doctors provide evidence-based care for patients with appetite problems and cachexia-anorexia syndrome.

The [eTG Palliative Care](#)³, particularly 'Gastrointestinal Symptoms in Palliative Care', is another useful resource (available through [www.ciap.health.nsw.gov.au](#) for NSW Health employees).

General information

There is limited evidence for efficacy of cannabis medicines in improving anorexia-cachexia related to cancer.

Prior to considering cannabis medicines as a treatment option for cachexia/anorexia syndrome in patients with advanced cancer, medical practitioners should be aware that there is limited evidence for efficacy

of cannabis medicines in this syndrome.

Medical practitioners should review the following guidance advice on made available by the [Therapeutic Goods Administration](#) (TGA) before making a decision to prescribe medicinal cannabis products:

- [Guidance for the use of medicinal cannabis in Australia: Overview](#)
- [Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia](#)

Summary of best practice in management of cachexia/anorexia syndrome in patients with advanced cancer

The following information is based on information from both the [Caresearch](#) website and [eTG Complete](#). More detailed information and prescribing guidelines regarding dose, frequency and sequential drug choices should be sought from these sources.

- Anorexia (lack of appetite) and weight loss are among the most recognised and troubling symptoms for patients, their family and carers.
- Cachexia is caused by tumour-related cytokines which can lead to catabolism with alteration in carbohydrate, protein and lipid metabolism.
- Preparing and eating food has social and emotional meaning for a patient and family.
- There is often a strong community belief that weight stabilisation postpones death. For most life-limiting illnesses (particularly advanced cancer) this is not the case. However palliative care has been shown to prolong survival.
- Potentially reversible and common contributors to appetite problems should be sought and treated as appropriate: mouth problems (thrush, mucositis), nausea, dysphagia, constipation.
- A focus on weight gain on its own may not result in meaningful clinical changes for palliative care patients. Pharmacotherapy should only be initiated for quality of life purposes.

Management


- Routine measuring of weight is not helpful and may contribute to patient and carer anxiety.
- Manage family, social and cultural expectations through compassionate discussion of expectations and outcomes.

- The prescriber should undertake a careful history and examination to reveal reversible and/or treatable factors.
- Non-pharmacological options include provision of small food portions (these are often better tolerated), focus on enjoyment of food rather than nutritional benefit and a strong focus on patient preference.
- Oral care should be provided and treatment initiated for oral thrush or mucositis (where present).
- Nausea, constipation and pain should be managed according to best evidence.
- Dysphagia should be investigated if appropriate. If a patient who has dysphagia wants to eat, consider referral to a dietician.
- Where depression is a contributing factor, consider whether an antidepressant is indicated. Mirtazapine may have beneficial ef-

fect on appetite.

- If slow gastric emptying is contributing to early satiety, commence metoclopramide or domperidone unless contraindicated.
- Corticosteroids can be considered. Careful consideration should be made to the risk vs. benefit profile and addition of a proton pump inhibitor. If loss of appetite is causing an impact on patient quality of life, a short-term (less than 14 days) trial of 2-4mg daily dexamethasone may be beneficial. If there is no improvement at day 5, cease therapy.
- There is evidence to support the use of either progestogens or corticosteroids as appetite stimulants in advanced cancer, but less evidence to suggest that they are associated with any improvement in quality of life.

Best Practice Guidance for Management of Anorexia and Cachexia in Palliative Care Patients with Advanced Cancer



Prescribing medicinal cannabis for the management of cachexia/anorexia syndrome in patients with advanced cancer

If the decision to proceed to cannabis prescription is made, please consider the following information.

Working through the following information will assist medical practitioners in completing the regulatory paperwork required to prescribe medicinal cannabis.

Further advice and assistance is available to medical practitioners via the [NSW Cannabis Medicines Advisory Service](#) by email or by telephone (02) 4923 6200.

The Prescriber

The prescriber should have a key involvement in provision of care. The prescriber may be a palliative care physician, an oncologist or a general practitioner in liaison with a specialist practitioner. Where the GP has a particular qualification in palliative care (e.g. Clinical Diploma in Palliative Care) it may be useful to provide this information. It is important that all members of the treating team are aware of the decision to prescribe a cannabis medicine – it is important that this information is communicated to other members of the care team in writing.

The prescriber should have had or will have an ongoing therapeutic relationship with the patient. Follow up and assessment of efficacy is essential.

Given the current status of cannabis medicines as unregistered medicines, prescribers should consider seeking advice from their medical indemnity insurers prior to prescribing.

The Patient

The patient must give informed consent to treatment. The consent process should include:

- This is an unregistered medicine (and costs may reflect this).
- The efficacy and side effects of this therapy is still being researched.
- However, they must be made aware of the likely effects and side effects of treatment.
- There will be restrictions on driving and operating heavy machinery.
- The patient should also be given clear information about therapeutic goals and likely stopping criteria.
- Potential for dependence or withdrawal

Patients should be advised that they are not able to drive while treated with medicinal cannabis.

Patients should be informed that measurable concentrations of THC can be detected in saliva for significant periods of time after administration. Further information is available from the [Transport for NSW Centre for Road Safety](#) and in [NSW Health's Prescribed Cannabis Medicines and Fitness to Drive Factsheet](#).

Prescribing a cannabis medicine: important considerations

The following has been adapted from the TGA's [Guidance for the use of medicinal cannabis in Australia: Patient information](#).

Cannabis products

- A variety of products are available.
- There are up to 100 cannabinoids (chemical compounds) in the cannabis plant.
- Many of the studies described in the medical literature have used either smoked cannabis (which is not recommended on health grounds) or purified tetrahydrocannabinol (THC) or cannabidiol (CBD).

- Tetrahydrocannabinol (THC) is responsible for the psychoactive effects of cannabis and is the reason cannabis is used recreationally. THC may contribute to reduction of nausea, vomiting, pain and muscle spasms as well as improvements in sleep and appetite.
- Cannabidiol (CBD) is not psychoactive and may be useful in the management of seizures, pain, and may have anxiolytic and anti-psychotic effects. Adding CBD to a THC product in a patient with toxicity to reduce toxicity is unproven. More appropriate is to reduce the dose and/or frequency of THC.
- Different cannabis products contain different ratios of THC to CBD.
- There are other cannabinoids under research including cannabigerol (CBG), tetrahydrocannabivarin (THCV), cannabinol (CBN) and cannabichromene (CBC).

Route of administration

Smoking

- Rapid onset of action, usually within minutes
- High levels of blood THC with shorter duration of effect
- Peak concentration in 30 minutes, effects last 2-4 hours
- Significant individual variability due to unknown quantity of THC in product and loss of active agent in side stream/combustion during smoking
- Smoking is well known to be harmful and not recommended on health grounds

Vaporising/oro-mucosal/buccal routes

- Rapid absorption and high blood concentrations
- Fewer toxins and reduced "side stream"
- Peak concentration in 15-30 minutes, effects last 2-4 hours
- Useful for symptoms requiring rapid and intermittent relief

Oral administration

- Oils or liquid capsules are available
- Slow onset of action with first effects at 30-90 minutes, peak effect at 2-4 hours
- Effects can last 8-24 hours
- Useful for symptoms requiring relief over longer periods of time (similar to controlled release medications)
- Titration of dosing may be easier with oro-mucosal sprays than oral formulations
- Sprays may be easier for those with difficulty swallowing

Topical

- Cannabinoids are lipophilic and therefore are likely to be well absorbed across the skin.
- THC is relatively less well absorbed than cannabidiol and cannabinol.
- Time of onset, duration of action and likelihood of depot occurrence are unknown and this route is therefore not recommended at this stage.

Evidence for Use

The following has been adapted from the TGA's [Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia](#)¹.

Five studies examined the use of medicinal cannabis for symptom control in patients with advanced cancer. Three studies were of moderate quality, and two of low quality. In a meta-analysis, there were no significant differences in outcomes between patients who received medicinal cannabis or placebo.

Thus, considering the evidence, a **THC-only product could be considered as the first cannabinoid to be added to a patient's treatment as this is where the best evidence of effect lies** (although this is small), noting that there is also evidence of potential harm (due

to side-effects) and lack of efficacy. Where a doctor chooses to use an alternative product to THC-only products, good clinical reasoning must be given, giving weight to the evidence above and understanding of cannabinoid pharmacology.

Evidence Grade	Cannabinoid used	Outcomes
C - Unclear scientific evidence for this use	Dronabinol, THC:CBD, THC	There was a non-significant effect of medicinal cannabis to reduce cancer pain.
D - Fair scientific evidence against this use (it may not work)	Dronabinol, THC:CBD, THC	There were no significant differences between medicinal cannabis and placebo in effects on patients' weight, caloric intake, appetite, nausea and vomiting, sleep, depressed mood, or quality of life. There were also no significant differences between medicinal cannabis and placebo in development of cognitive impairment or dizziness during treatment.
C - Unclear scientific evidence for this use	Cannabis sativa	All cancer and anti-cancer treatment symptoms (e.g. nausea, vomiting, mood disorders, fatigue, weight loss) were improved after initiating cannabis therapy. The lack of controlled studies restricts the conclusiveness of this finding and the clinical relevance is also unknown.
C - Unclear scientific evidence for this use	Dronabinol	Patients receiving dronabinol reported an increase in appetite and a decrease in nausea, however the treatment effects decreased as the disease progressed. The lack of controlled studies restricts the conclusiveness of this finding.
C - Unclear scientific evidence for this use	Nabilone	Patients receiving nabilone reported decreases in pain scores, morphine use, nausea, anxiety, overall distress, and borderline significant improvement in appetite. The lack of controlled studies restricts the conclusiveness of this finding.

Table 1 —Extract from *Symptom Control in Cancer*, in [Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia](#)¹. For additional information visit the TGA's [Summary of Evidence](#) [here](#).

Precautions

The following has been adapted from the TGA's [Guidance for the use of medicinal cannabis in Australia: Overview](#)⁴.

Medicinal cannabis products containing THC are generally **not appropriate** for patients who:

- Have a previous psychotic or concurrent active mood or anxiety disorder.
- Are pregnant, planning on becoming pregnant, or breastfeeding.
- Have unstable cardiovascular disease.

Any allergies to potential carrier oils (e.g. sesame, canola, sunflower) should be noted as products available in Australia often contain oils. This may influence product selection.

Consider concomitant medication use and potential for pharmacokinetic and pharmacodynamic drug interactions (see below).

Adverse effects

The following has been adapted from the TGA's [Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia](#)¹.

Patients and prescribing clinicians should be aware of possible adverse events such as somnolence, nausea and dizziness. Adverse events such as confusion, pain, diarrhoea or hallucinations may impact the overall aims of the palliative medicine and reduce quality of life, and should be evaluated on a case-by-case basis.

Serious adverse events (SAEs) have been described clinically, such as psychosis and cognitive distortion requiring external assistance, but the severity, time course and further details are not recorded. SAEs are also likely where there are concomitant medications prescribed that are metabolised by, induced or inhibited by enzymes in the P450 system, or where there are similar pharmacodynamic effects.

Based on the available studies, commonly reported adverse events in the use of medicinal cannabis products in the palliative care setting include:

- nausea (21% of patients)
- somnolence (20% of patients)
- dizziness (16% of patients)
- asthenia (13% of patients)
- tiredness/fatigue (12% of patients)
- vomiting (11% of patients)
- anaemia (11% of patients)
- confusion (10% of patients)
- pain (10% of patients)
- diarrhoea (8% of patients)
- headache (8% of patients)
- dyspnoea (8% of patients)
- hallucinations (5% of patients)
- In a small (15 patient) study, 11 had anxiety symptoms.

Drug Interactions

The current available information on drug interactions with cannabinoids is mainly sourced from the product information on Marinol®⁵ (THC) and Sativex®⁶ (THC and CBD combination). Drug interactions with cannabidiol have been summarised in the literature^{7,8}.

Ongoing reports of potential interactions to the TGA is vital to improve the data in this area.

Pharmacokinetic interactions

- delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) are metabolised by the cytochrome P450 enzyme system. CBD has significant inhibitory effect on this system, in addition.
- In vitro studies have shown Sativex® has CYP450 enzyme inhibition effects.
- If concomitant drug treatment with CYP3A4 inhibitors (e.g. ketoconazole, ritonavir, clarithromycin, fluoxetine) is started or stopped, a new dose titration may be required. If concomitant drug treatment with strong enzyme inducers (e.g. rifampicin, carbamazepine, St John's Wort) is started or stopped, a new dose titration

may be required.

- Monitor for potentially increased delta-9-THC -related adverse reactions when co-administered with inhibitors of CYP2C9 (e.g., amiodarone, fluconazole) and inhibitors of CYP3A4 enzymes (e.g., ketoconazole, itraconazole, clarithromycin, ritonavir, erythromycin, grapefruit juice).
- delta-9-THC is highly bound to plasma proteins and might displace and increase the free fraction of other concomitantly administered protein-bound drugs. Although this displacement has not been confirmed *in vivo*, monitor patients for increased adverse reactions to narrow therapeutic index drugs that are highly protein-bound (e.g. warfarin, cyclosporine, amphotericin B) when initiating treatment or increasing the dosage of delta-9-THC
- Based on *in vitro* data an inhibition of p-glycoprotein at the intestinal level by CBD cannot be excluded. Therefore, caution is recommended upon concomitant treatment with digoxin and other substrates for p-glycoprotein.

Pharmacodynamic interactions

Sedation

- Care should be taken with hypnotics, sedatives and drugs with potential sedating effects as there may be an additive effect on sedation and muscle relaxing effects. This may include opioids, benzodiazepines, anticholinergics and antihistamines.
- Cannabis medications may interact with alcohol, affecting co-ordination, concentration and ability to respond quickly. In general, alcoholic beverages should be avoided

Cardiac toxicity

- Additive hypertension, tachycardia, possibly cardiotoxicity with amphetamines, cocaine, atropine or other sympathomimetic agents.

Psychiatric effects

- Case reports of hypomania when cannabis co-administered with fluoxetine and also disulfiram.

Musculoskeletal

- Care should be taken when co-administering Sativex with anti-spasticity agents since a reduction in muscle tone and power may occur, leading to a greater risk of falls.

Dosing

Prescribers who are unfamiliar with these products should consider contacting the [NSW Cannabis Medicines Advisory Service](#) by [email](#) or by telephone (02) 4923 6200.
In general:

- Start at low dose and frequency.
- Titrate to effect whilst monitoring for side effects.
- Patient response to these medications varies widely.
- Favour lower doses in the elderly who may be at higher risk of CNS and cardiac adverse effects.

The following are examples only – doses should be individualised for patient and indication.

Example for THC – adapted from Dronabinol (Marinol®) product information⁵ (for treatment of anorexia):

- Starting dose 2.5mg once daily or BD
- Monitor for side effects
- If tolerated, and further therapeutic effect desired, slowly increase dose.
- Higher doses are variably tolerated:
 - 10 mg twice daily has been tolerated in about half of the patients in appetite stimulation studies. It is likely that there will be less tolerance in older patients or those with significant physiologic compromise.
 - Generally there should be a maximum dose of 30mg of tetra-

hydrocannabinol per day. Beyond this dose, the risk of adverse effects may increase.

Example for THC:CBD – from Sativex® oromucosal spray Product Information⁶. Note that this PI was developed for the indication of spasticity in multiple sclerosis.

- Initially one spray per day, slow titration over two weeks.
- The mean number of sprays required for symptom relief was 4.81 per day (i.e. THC 12.5mg/CBD 12.98mg in divided doses daily).

Monitoring Outcomes

There are three areas of outcomes that should be considered:

1. Symptom control

A pre-defined measure of success should be negotiated with the patient prior to commencement of therapy. This can be measured by using a validated tool, for example by the [Palliative Care Symptom Assessment Scale](#).

2. Drug adverse events

Careful monitoring of patients for adverse events and the need for a change in dosage is important. Adverse events may become apparent after commencement or after change in dose. Adverse events may be related to other concurrent medications. Doses of these medicines should be adjusted as appropriate. Significant adverse events **must** be reported to the TGA (including dependence and withdrawal symptoms)

3. Pathology monitoring

Drug concentrations of drugs where there is a potential or actual drug-drug interaction may be important.

NEED FURTHER INFORMATION AND ASSISTANCE?

Further advice and assistance is available to all NSW-based medical practitioners through the [NSW Cannabis Medicines Advisory Service](#).

The Service can be contacted by [email](#) or by telephone (02) 4923 6200.

References

1. Department of Health Therapeutic Drug Administration. Guidance for the use of medicinal cannabis in the treatment of palliative care patients in Australia, December 2017. <https://www.tga.gov.au/sites/default/files/guidance-use-medicinal-cannabis-treatment-palliative-care-patients-australia.pdf>
2. Caresearch – the Palliative Care Knowledge Network <https://www.caresearch.com.au/caresearch/tabid/182/Default.aspx> (accessed 28 Feb 2018)
3. eTG Complete <https://tgldcdp.tg.org.au/etgcomplete> (accessed 28 Feb 2018)
4. Department of Health Therapeutic Drug Administration. Guidance for the use of medicinal cannabis in Australia: Overview. <https://www.tga.gov.au/publication/guidance-use-medicinal-cannabis-australia-overview>.
5. Marinol Product Information https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/018651s029lbl.pdf (accessed 28 Feb 2018)
6. Sativex Product Information <https://www.tga.gov.au/sites/default/files/auspar-nabiximols-130927-pi.pdf> (accessed 28 Feb 2018)
7. Sarah Melton. Stirring the Pot: Potential Drug Interactions With Marijuana - Medscape - Jun 08, 2017 <https://www.medscape.com/viewarticle/881059>.
8. Gaston, T. E., Bebin, E. M., Cutter, G. R., Liu, Y., Szaflarski, J. P. and the UAB CBD Program (2017), Interactions between cannabidiol and commonly used antiepileptic drugs. *Epilepsia*, 58: 1586–1592. doi:10.1111/epi.13852.